

Références

- Bennett, A. L., Lavie, C. J., & Grace, S. L. (2017). Cardiac Rehabilitation Following Acute Coronary Syndrome in Women. *Curr Treat Options Cardiovasc Med*, 19(8), 57. doi:10.1007/s11936-017-0559-x
- Dreyer RP, Sciria C, Spatz ES, Safdar B, D'Onofrio G, Krumholz HM. Young women with acute myocardial infarction: current perspectives. *Circ Cardiovasc Qual Outcomes*. 2017;10(2)
- Feola M, Garnero S, Daniele B, et al. Gender differences in the efficacy of cardiovascular rehabilitation in patients after cardiac surgery procedures. *J Geriatr Cardiol*. 2015;12(5):575–9
- Halm, M. and Penque, S. 1999. Heart disease in women. *American Journal of Nursing*, 99(4y.26-3i.
- Hart PL. Women's perceptions of coronary heart disease: an integrative review. *J Cardiovasc Nurs* 2005; 20: 170–6.
- Kristofferzon, M. L., Lofmark, R., & Carlsson, M. (2007). Striving for balance in daily life: experiences of Swedish women and men shortly after a myocardial infarction. *J Clin Nurs*, 16(2), 391-401. doi:10.1111/j.1365-2702.2005.01518.x
- Oliver-McNeil S, Artinian NT. Women's perceptions of personal cardiovascular risk and their risk-reducing behaviours. *Am J Crit Care* 2002;11(3):221 – 7.
- Vaccarino, V., Krumholz, H. M., Yarzebski, J., Gore, J. M., & Goldberg, R. J. (2001). Sex differences in 2-year mortality after hospital discharge for myocardial infarction. *Annals of Internal Medicine*, 134, 173–181.
- Wenger NK. Coronary heart disease: the female heart is vulnerable. *Prog Cardiovasc Dis* 2003; 46: 199–229.